

## Fracture of the base of the 5<sup>th</sup> metatarsal (Jones)

Foot 2

### Fracture Care Team: Shared Care Plan

Trust Switchboard for both sites: 0300 131 4500

Fracture clinic and orthopaedic outpatient appointments:

Eastbourne 0300 131 4788

Conquest 0300 131 4861

Casting Department: Eastbourne 0300 131 5564

Casting Department: Conquest 0300 131 4860

**This information leaflet explains the ongoing management of your injury.**


**You have sustained a fracture to the base of the 5<sup>th</sup> metatarsal of your foot, which is known as a Jones fracture. Please use the picture below to understand where this injury is.** This normally takes approximately 6 weeks to unite (heal) although pain and swelling can be ongoing for 3-6 months. You will be in a cast and asked to keep your weight off the foot initially. The swelling is often worse at the end of the day and elevating it will help. Following fracture clinic review you may be provided with a boot instead of the cast and you will be advised on how much weight to put through the foot. The boot which you may have been given is for your comfort only and is not needed to aid fracture healing. Take pain killers as prescribed. If you are worried that you are unable to follow this rehabilitation plan, or have any questions, then please phone the Fracture Care Team for advice.

If you are experiencing pain or symptoms, other than at the site of the original injury or surrounding area, please get in touch using the telephone number at the top of this leaflet.

You have also been referred to fracture clinic after your injury to check on your progress. They will offer on-going advice with regards to rehabilitation. They may require further x-rays and this will be decided by the consultant on the day. There is a small chance that the fracture does not heal naturally and surgery may need to be considered. This will be discussed in the consultation at your appointment.

If you have not received an appointment within one week please contact the Fracture Care Team on the details provided above.

### Picture of injury:



## Please follow the Management / Rehabilitation plan shown below:

Weeks since injury	Rehabilitation plan
0- 6	While in the cast, keep your weight off the foot using the crutches. Once reviewed in clinic you may be changed into a black boot which you must wear all of the time when walking. Use the crutches to take some of the weight off your foot. It is ok to take the boot off at night and when resting at home. It is also important to perform the exercises below regularly to get the movement back.
6 -12	You have been reviewed in fracture clinic. If you have been advised that the fracture is united (healed), you can begin to resume normal activity but be guided by any pain you are experiencing. You should be able to carry out day to day activities. Arduous tasks, long walks etc., may still cause some discomfort and swelling.

### Initial advice

#### Cold packs:

A cold pack (ice pack or frozen peas wrapped in a damp towel) can provide short term pain relief. Apply this to the sore area for up to 15 minutes, every few hours ensuring the ice is never in direct contact with the skin.

#### Rest and Elevation:

Try to rest the foot for the first 24-72 hours to allow the early stage of healing to begin. Raise your ankle above the level of your hips to reduce swelling. You can use pillows or a stool to keep your foot up.

#### Early movement and exercise:

Early movement of the ankle and foot is important to promote circulation and reduce the risk of developing a DVT (blood clot). Follow the exercises below without causing too much pain. This will ensure your ankle and foot do not become too stiff. These exercises will help the healing process.

Early weight bearing (putting weight through your injured foot) helps increase the speed of healing. Try to walk as normally as possible as this will help with your recovery.

### Smoking cessation

Medical evidence suggests that smoking prolongs fracture healing time. In extreme cases it can stop healing altogether. It is important that you consider this information with relation to your recent injury. Stopping smoking during the healing phase of your fracture will help ensure optimal recovery from this injury.

For advice on smoking cessation and local support available, please refer to the following website: <http://smokefree.nhs.uk> or discuss this with your GP.

## Initial exercises (3 to 4 times a day)

**Ankle and foot range of movement exercises. Repeat these 10 times each.**

1. Point your foot up and down within a comfortable range of movement.
2. Make circles with your foot in one direction and then change direction.
3. With your heels together, move your toes apart, as shown in the picture.



## Sources of information

This information leaflet has been developed by the Fracture Care Team at Brighton and Sussex University Hospitals Fracture Care Team and adapted for use at East Sussex Healthcare NHS Trust. Information updated during the COVID-19 pandemic to ensure that patients with injuries have information, support and care despite social distancing.

## Important information

This patient information is for guidance purposes only and is not provided to replace professional clinical advice from a qualified practitioner.

## Your comments

We are always interested to hear your views about our leaflets. If you have any comments, please contact the Patient Experience Team – Tel: 0300 131 4731 (direct dial) or by email at: [esh-tr.patientexperience@nhs.net](mailto:esh-tr.patientexperience@nhs.net)

## Hand hygiene

The trust is committed to maintaining a clean, safe environment. Hand hygiene is very important in controlling infection. Alcohol gel is widely available at the patient bedside for staff use and at the entrance of each clinical area for visitors to clean their hands before and after entering.

## Other formats

**If you require any of the Trust leaflets in alternative formats, such as large print or alternative languages, please contact the Equality and Human Rights Department.**

**Tel: 0300 131 4500 - Email: [esh-tr.AccessibleInformation@nhs.net](mailto:esh-tr.AccessibleInformation@nhs.net)**

After reading this information are there any questions you would like to ask? Please list below and ask your nurse, practitioner or doctor.

---

---

---

---

---

## Reference

Thank you to the Brighton and Sussex University Hospitals NHS Trust and to Kate Weatherly, Specialist Orthopaedic Physiotherapist for the first version.

The following clinicians have been consulted and agreed this patient information:

- Mr Guy Selmon – Consultant Orthopaedic Surgeon
- Mr Andrew Skyrme – Consultant Orthopaedic Surgeon
- Mr Michael Dunning - Consultant Orthopaedic Surgeon
- Mr Henry Willmott - Consultant Orthopaedic Surgeon
- Mr Barry Rose - Consultant Orthopaedic Surgeon
- Helen Harper-Smith – Professional Lead Physiotherapist
- Hilary Kircher – Clinical / Operational Lead Outpatient Physiotherapist

The Clinical Specialty/Unit that have agreed this patient information leaflet:  
Department of Trauma and Orthopaedics

Next review date: March 2023  
Responsible clinician/author: Helen Harper-Smith– Professional Lead Physiotherapist  
Mrs Scarlett McNally – Consultant Orthopaedic Surgeon

This leaflet can also be found at: [www.esht.nhs.uk/leaflet/fracture-of-the-base-of-the-5th-metatarsal-jones/](http://www.esht.nhs.uk/leaflet/fracture-of-the-base-of-the-5th-metatarsal-jones/)

© East Sussex Healthcare NHS Trust – [www.esht.nhs.uk](http://www.esht.nhs.uk)