

HOW TO TAKE PLENVU®



AFTERNOON APPOINTMENT

For full details, please refer to the patient information leaflet (PIL) supplied with your box of Plenvu.

Reporting of side effects in the United Kingdom

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard.

By reporting side effects you can help provide more information on the safety of this medicine.

Norgine Adverse events should also be reported to Medical Information at Norgine Pharmaceuticals on +44 1895 826606 or E-mail: medinfo@norgine.com


Reporting of side effects in Ireland

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly to the HPRA, via www.hpra.ie, medsafety@hpra.ie or by calling (01) 676 4971. By reporting side effects you can help provide more information on the safety of this medicine.

Norgine Adverse events should also be reported to Medical Information at Norgine Pharmaceuticals on +44 1895 826606 or E-mail: medinfo@norgine.com

CLEAR FLUIDS

Remember you can keep drinking until 2 hours before your colonoscopy.



EATING SUGGESTIONS

Your healthcare professional may request that you adopt a low fibre diet when preparing for your bowel preparation.

YES ✓ LOW FIBRE FOODS WHICH ARE PREFERRED



NO ✗ HIGH FIBRE FOODS WHICH SHOULD BE AVOIDED



Supplied as a service to patients by Norgine Pharmaceuticals Limited.
PLENVU, NORGINE and the sail logo are registered trademarks of the Norgine group of companies
UKE-COR-PR-2000012 Date of preparation: April 2020



Unless your doctor or nurse tells you differently, start taking Plenvu as outlined below.

For full details, please refer to the patient information leaflet (PIL) supplied with your box of Plenvu.


All sachets must be taken with the right amount of water according to the timings below.

You will have lots of watery bowel movement, so **stay near a toilet** because you may need to use it urgently.

DAY BEFORE colonoscopy



DAY OF colonoscopy



DOSE 1 6pm



Open the carton and remove the **Dose 1** sachet



Pour the contents of the **Dose 1** sachet into a measuring container that can hold at least 500ml of fluid



Add water to make up to 500ml and stir until all the powder has fully dissolved



Each dose of Plenvu must be **sipped slowly** over 30 minutes, followed by 500ml of mandatory clear fluid over a further 30 minutes



DOSE 2 6am



Open the carton and remove the **Dose 2** sachets A and B



Pour the contents of **Dose 2** sachets A and B into a measuring container that can hold at least 500ml of fluid.



Add water to make up to 500ml and stir until all the powder has fully dissolved



Each dose of Plenvu must be **sipped slowly** over 30 minutes, followed by 500ml of mandatory clear fluid over a further 30 minutes.

Hints & Tips



Sip slowly over at least 30 minutes



Once dissolved, (this may take up to 8 minutes) the solution can be refrigerated



Try using a straw



Stay hydrated with plenty of clear fluids*

*Please see overleaf.