

## Minimally displaced glenoid fracture

### Fracture Care Team: Shared Care Plan

Trust Switchboard for both sites: 0300 131 4500

Fracture clinic and orthopaedic outpatient appointments:

Eastbourne 0300 131 4788


Conquest 0300 131 4861

This information leaflet explains the ongoing management of your injury.

**You have sustained a minimally displaced fracture to the Glenoid in your shoulder.** The shoulder is a ball and socket joint. You have fractured the socket part. This normally takes between 6-12 weeks to unite (heal). Once the fracture begins to heal it is important to keep the shoulder moving to overcome the stiffness but not to aggravate it. The shoulder joint does not respond well to being injured and longstanding stiffness is often inevitable.

Take pain killers as prescribed. You may find it more comfortable to sleep propped up with pillows. If you are worried that you are unable to follow this rehabilitation plan, or have any questions, then please contact the Fracture Care Team for advice.

### Picture of injury



Although the fracture is in a good position and only has a very small chance of moving, we routinely make an appointment in the fracture clinic 1-2 weeks after your injury. You will have another x-ray at this appointment. The Specialist will then assess your shoulder and guide the next stage of your rehabilitation.


If you have not received this appointment within one week of receipt of this letter please contact the Fracture Care Team on the number provided above.

## Please follow the Management/rehabilitation plan shown below

Weeks since injury	Rehabilitation plan
0 - 4	Wear the sling all the time, even in bed at night. Begin to do the Initial exercises shown below. You will be seen in fracture clinic.
4 - 6	Continue to wear the sling. Unless you have been advised otherwise, you may progress to the Stage 2 exercises. Do not lift your elbow above shoulder height as this may cause excessive pain.
6 - 12	You may start to discard the sling. Begin normal light activities with the arm and shoulder. Increase movement using the Stage 3 exercises. The fracture should be largely united (healed). You should be able to increase day to day activities. More arduous tasks may cause discomfort. Start to lift your arm overhead if possible.
12	<b>If you are still experiencing significant pain and stiffness then please contact us for further consultation</b>

## Initial Exercises to do 4 – 5 times a day

If you have stiffness in your elbow or hand from wearing the sling, you may wish to perform these exercises first. However, once they become easy you can start with the posture and pendulum exercises.


**Elbow Bend to Straighten**

Bend and straighten your elbow so you feel a mild to moderate stretch. You can use your other arm to assist if necessary. Do not push into pain.

**Forearm Rotations**

Put your elbow at your side. Bend it to 90 degrees. Slowly rotate your palm up and down until you feel a mild to moderate stretch. You can use your other arm to assist if necessary. Do not push into pain.

Repeat 10-15 times provided there is no increase in symptoms.



### **Finger and wrist flexion and extension**

Open and close your hand as shown 10-15 times. Then move your wrist up and down 10-15 times.

After a few days, hold a soft ball/ball of socks. Squeeze the ball as hard as possible without pain. Hold for 5 seconds and repeat 10 times.



### **Postural awareness**

Bring your shoulders back and squeeze your shoulder blades together as shown in the picture. Do this with or without your sling on.

Hold the position for 20-30 seconds and repeat 5 times provided there is no increase in symptoms.



### **Shoulder pendular exercises**

Stand and lean forward supporting your injured arm with your other hand as shown in the picture. Try to relax your injured arm.

1. Assist your arm slowly and gently forwards and backwards.
2. Assist your arm slowly and gently side to side.

Continue for approximately 1-2 minutes in total provided there is no increase in symptoms. Remember to try and relax your arm.

## Stage 2 exercises to do 4-5 times a day - To start at 4 weeks



shown in the pictures.

### **Active assisted Shoulder flexion**


Use your other hand to lift your arm up in front of you as

Repeat 10 times provided there is no increase in symptoms.



### **Active-assisted Abduction:**

Hold a stick in both hands as in the photo. Gently push your injured arm to the side (away from your body) as far as comfort allows. It may help to face a mirror initially to make sure the top of your shoulders stay level.



### **Active assisted External rotation**

Keep the elbow of your injured arm tucked into your side and your elbow bent. Hold onto a stick/umbrella/golf club or similar. Use your unaffected arm to push your injured hand outwards. Remember to keep your elbow tucked in. Push until you feel a stretch.

If you don't have a stick you could simply hold the injured arm at the wrist and guide it outwards. Hold for 5 seconds then return to the starting position. Repeat 10 times provided there is no increase in symptoms.

## Stage 3 exercises to do 4-5 times a day - To start at 6 weeks

When you have regained full range of movement during the above exercises without pain you can start to do the exercises without the support of your other hand; this is known as active range of movement. Then when you have regained your full range of movement without the support of the other arm you can start to build up your regular activities.




### **Active Forward flexion:**

With your thumb facing up, try to move your arm up, keeping it close beside your body.

### **Active Abduction**

With your thumb facing up and outwards, try to move your arm in a big arc out to the side.



### **Active External rotation**

With your elbow by your side, rotate your forearm outwards, keeping your elbow at about 90 degrees in flexion.



Repeat all of these 3 exercises 10 times each, 4-5 times a day. Only go as far as you can naturally, without doing any trick movements to try and get further. This will increase over time and should not be forced.

If you are having problems progressing with the exercises and have a follow-up consultation booked, please do let the clinician know so that they can review the exercises and refer you on to Physiotherapy if necessary. If you are on an independent management programme, then please contact us using the number at the top of the letter so that we can also arrange physiotherapy for you.

### **Smoking cessation**

Medical evidence suggests that smoking prolongs fracture healing time. In extreme cases it can stop healing altogether. It is important that you consider this information with relation to your recent injury. Stopping smoking during the healing phase of your fracture will help ensure optimal recovery from this injury.

For advice on smoking cessation and local support available, please refer to the following website: <http://smokefree.nhs.uk> or discuss this with your GP.

### **Sources of information**

This information leaflet has been developed by the Fracture Care Team at Brighton and Sussex University Hospitals Fracture Care Team and adapted for use at East Sussex Healthcare NHS Trust. Information updated during the COVID-19 pandemic to ensure that patients with injuries have information, support and care despite social distancing.

### Important information

This patient information is for guidance purposes only and is not provided to replace professional clinical advice from a qualified practitioner.

### Your comments

We are always interested to hear your views about our leaflets. If you have any comments, please contact the Patient Experience Team – Tel: 0300 131 4731 or by email at: [esh-tr.patientexperience@nhs.net](mailto:esh-tr.patientexperience@nhs.net)

### Hand hygiene

The trust is committed to maintaining a clean, safe environment. Hand hygiene is very important in controlling infection. Alcohol gel is widely available at the patient bedside for staff use and at the entrance of each clinical area for visitors to clean their hands before and after entering.

### Other formats

**If you require any of the Trust leaflets in alternative formats, such as large print or alternative languages, please contact the Equality and Human Rights Department.**

**Tel: 0300 131 4434 Email: [esh-tr.AccessibleInformation@nhs.net](mailto:esh-tr.AccessibleInformation@nhs.net)**

After reading this information are there any questions you would like to ask? Please list below and ask your nurse, practitioner or doctor.

---

---

---

---

### Reference

The following clinicians have been consulted and agreed this patient information:

Dr Danielle Vidler – Consultant and Clinical lead Emergency Department CQ

Mr Utham Shanker – Consultant and Clinical lead Emergency Department EDGH

The directorate group that has agreed this patient information leaflet:

URGENT CARE

Next review date: December 2027

Responsible clinician/author: Emergency Department Consultants

© East Sussex Healthcare NHS Trust – [www.esht.nhs.uk](http://www.esht.nhs.uk)